

MAGNETIC ORIENTATION IN ANURANS: A CASE STUDY ON *RHINELLA ARENARUM* (ANURA: BUFONIDAE)

ORIENTACIÓN MAGNÉTICA EN ANUROS: UN ESTUDIO DE CASO SOBRE *RHINELLA ARENARUM* (ANURA: BUFONIDAE)

Mario Andrés Herrera Moratta^{*}, Ana Laura Navas Romero^{1,2}, Lorena Quiroga³, Consuelo Escudero⁴, Eduardo Sanabria³

¹Instituto Argentino de Investigaciones en Zonas Áridas-CONICET-Mendoza, Argentina. CP 5500, Argentina.

²Instituto de Ingeniería Química-Facultad de Ingeniería (UNSJ)-Grupo Vinculado al PROBIEN (CONICET-UNCuyo), San Juan, CP 5400, Argentina.

³Instituto de Ciencias Básicas. Facultad de Filosofía, Humanidades y Arte. Universidad Nacional de San Juan. CONICET. San Juan, CP 5400, Argentina.

⁴Facultad de Ciencias Exactas, Físicas y Naturales. Universidad Nacional de San Juan. San Juan, CP 5400, Argentina.

*Correspondence: maherrera@mendoza-conicet.gob.ar

Received: 2023-03-29. Accepted: 2023-08-22. Published: 2023-09-05.

Editor: Adriana Manzano, Argentina.

Resumen.— El campo magnético de la Tierra (EMF) es una de las fuentes de información direccional más uniformes y accesibles que los animales pueden usar en los movimientos de comportamiento. El presente estudio tuvo como objetivo determinar la capacidad de *Rhinella arenarum* para percibir los campos electromagnéticos. Para evaluar la influencia de los campos electromagnéticos en los anuros recolectamos cuarenta y cuatro especímenes adultos de *R. arenarum* durante la temporada de reproducción. Registramos el movimiento de cada sapo desde el centro hasta la periferia de un arenero circular utilizando una cámara de visión nocturna. Repetimos el experimento después de cinco minutos, con un campo magnético inducido adicional (IMF), que fue creado empleando dos bobinas de aire Helmholtz. Los movimientos de los sapos bajo la presencia de EMF y IMF fueron significativamente diferentes. Concluimos que *R. arenarum* podría usar el EMF como mecanismo de navegación y sistema de ubicación, para viajar largas distancias hasta los estanques de desove año tras año.

Palabras claves.— Navegación, migración de anuros, percepción magnética, ubicación.

Abstract.— The Earth's Magnetic Field (EMF) is one of the most uniform and accessible sources of directional information that animals can use for behavioral movements. The present study aimed to determine the ability of *Rhinella arenarum* to perceive the EMF. To assess the influence of the EMF on anurans *in situ* we collected forty-four adult specimens of *R. arenarum* during the breeding season. We recorded the movement of each toad from the center to the periphery of the circular arena utilizing a night vision camera. We repeated the experiment after five minutes, with an additionally Induced Magnetic Field (IMF), which was created by two Helmholtz air coils. The movements of the toads under the presence of the EMF and IMF were significantly different. We concluded that *R. arenarum* could use the EMF as a navigation mechanism and location system, to travel long distances to spawning ponds year after year.

Key Words.— Navigation, anuran migration, magnetic perception, location.

INTRODUCTION

Animals exploit numerous sources of information while migrating, homing, or moving around their habitats (Freake et al., 2005; Lohmann et al., 2007; Shakhparonov et al., 2022). Among these, the Earth's magnetic field (EMF) is a particularly

pervasive environmental feature (Skiles, 1985). In contrast to other cues, the magnetic field is present both day and night, is mostly unaffected by weather and seasonality, and exists everywhere on the planet (Lohmann et al., 2007; Wiltschko &

Wiltschko, 2012). Some species can find their way back home after being displaced to an unknown location, despite the absence of geographical references, goal-emanating cues, and directional information obtained during the journey (Phillips, 1996).

Homing behavior in animals is defined as an individual's ability to navigate through a small or unfamiliar terrain back to its "home" place, its home range (Nothacker et al., 2018). Various advantages of such site fidelity have been discussed, particularly the maximization of resource exploitation, which is often linked to reproduction (Nothacker et al., 2018). This ability to navigate requires, in addition to a directional sense (compass), an understanding of the geographical position by processing the available spatial information in the place where the individual is located (Diego-Rasilla & Luengo, 2020; Kramer, 1953). This phenomenon is found in many arthropods and vertebrates, such as decapod crustaceans (Pittman & McAlpine, 2003), salmon (Quinn & Dittman, 1990), or pigeons (Walcott, 1996). Among amphibians, it is known that many species exhibit site fidelity (Sinsch, 2014).

During the spawning season, several species of terrestrial anurans return to water bodies for mating mainly type I and II reproductive modes sensu Haddad & Prado (2005). In this process, the females release eggs that are fertilized by males (Young et al., 2004). As summarized by Wells (2007), the homing behavior of salamanders and newts is relatively well understood compared to other amphibians. Mechanisms or processes leading to the ability to return to the same spawning location year after year are still poorly unknown. Several studies show that amphibians are able to use a magnetic compass for orientation and navigation to the breeding ponds (Phillips et al., 1995; Diego-Rasilla, 2003; Diego-Rasilla et al., 2005, 2008, Sinsch, 2006, Phillips et al., 2022; Landler, 2022). The use of the geomagnetic field would be essential in situations where other positioning sources (maps) and directional cues (compasses), such as olfactory signals (and celestial cues are unavailable (Joly & Miaud, 1993; Diego-Rasilla et al., 2005). It has been demonstrated that magnetic field compasses are used in homing orientation by alpine newts (Diego-Rasilla, 2003) and in orientation towards the coast by eastern red-spotted newts (Phillips & Borland, 1992). Due to this, investigations on anuran migrations using the terrestrial magnetic field are important in order to clarify its effect. Studies on adult specimens of *Pelophylax ridibundus* showed the ability of anurans to use the Earth's magnetic field for orientation preference (Shakhparonov & Ogurtsov, 2017). In migratory studies of adult anurans, Landler & Gollmann (2011) made observations on *Bufo bufo* and its ability to use the magnetic field. After capturing specimens

migrating to spawning ponds, they tested their ability to correct their orientation toward the ponds in a circular arena in laboratory conditions. They demonstrated the influence of the Earth's Magnetic Field on their orientation behavior. However, the specimens were not orient themselves towards the spawning ponds but rather maintained their orientation prior to capture (orientation of the "d" axis; see Endler, 1970). Landler et al. (2016) repeated this experiment and tested *B. bufo* specimens the same night of capture, and after keeping them in captivity for three days, presumably enough time for them to update their internal map. The specimens tested on the same night of capture maintained the migration direction prior to capture, while the specimens kept in captivity exhibited random orientation. Extending their studies on *B. bufo*, in a new experiment Pail et al. (2020), examined the true navigation response in an open-air setting after keeping the toads in quarantine four days. In addition to the magnetic field, they measured the effect of environmental variables and the lunar cycle on the orientation of *B. bufo*. Their results showed that the animals were not yet oriented towards the breeding pond, but rather toward their previous migration direction prior to capture.

This investigation aimed, firstly, to determine whether males of *Rhinella arenarum* (Hensel, 1867) exhibit "homing" behavior toward the reproductive site from which they were displaced, and secondly, to determine the influence of the magnetic field on their orientation behavior towards the reproductive site in the arid region of San Juan, Argentina. We tested two hypothesis a) if we displace male frogs of *Rhinella arenarum* from their spawning pond, they will maintain orientation to their initial spawning site, and b) if we modify the Earth's Magnetic Field, adult males of *Rhinella arenarum* will change their orientation according to the new magnetic location of the spawning pond. The experiments presented in this study constitute a first approach to understand the orientation behavior in Neotropical anurans

MATERIALS AND METHODS

The experiments were carried out under natural climatic conditions after sunset at 8:30 p.m., from September 3rd to September 10th, 2012, in the Sarmiento Provincial Park. This park is located in Zonda, 25 km west of the city of San Juan (3° 33' 00.33" S, 68° 32' 22.58" W; elevation: 800 m a.s.l.). The area is characterized by the presence of extensive wetlands (40 ha), which are irrigated by the subterranean watercourse of the San Juan River. During the winter season, the wetland is reduced to two small water bodies, while the flooded areas expand during the summer, until they become a large flooded area (Victoria, 1999). Study area is located in Monte phytogeographical

province (Cabrera, 1971), a region characterized by an arid climate with a mean annual temperature of 17.3 °C, a mean annual maximum of 25.7 °C, and a mean annual minimum of 10.4 °C. Rainfalls mainly occur during the summer season, with a mean annual precipitation of 84 mm (Camarillo-Naranjo et al., 2018; Poblete & Hryciw, 2017).

The common toad (*R. arenarum*) is a widely distributed species in north and central Argentina inhabits various types of environments, including anthropogenic ones such as residential yards and gardens (Sotomayor et al., 2012; Sanchez et al., 2014). In the Monte Desert region of central Argentina, *R. arenarum* breeds from late August until late November (Sanabria et al., 2005). During the reproductive process, the females of this species deposit their eggs in gelatinous strands, laying an average of 28,000 eggs per clutch (Quiroga & Sanabria, 2012). There is sexual dimorphism in this species, with females being larger than males (Quiroga et al., 2004). As for its diet, it primarily feeds on Hymenoptera, specifically Formicidae (ants) (Quiroga, 2006). *Rhinella arenarum* is primarily a thermoconforming organism, with ambient temperature and wind speed being the environmental variables that most affect its body temperature (Sanabria et al., 2012).

We collected adult males (*R. arenarum*) from ponds located 500 m in the SE direction of the experimental area. Then, we transport the individuals to the experiment area in dark plastic containers to avoid any type of signal during the movement (0.3 m in diameter filled with 0.8-1.0 l of pond water).

We tested the behavior of the toads one by one on a circular arena 1.2 m in diameter, made with a medium-density fiberboard (MDF) wall 0.6 m high. We painted the wall as well as the bottom black color, so the toads inside of the sandbox could not distinguish the line from the horizon. We made two Helmholtz-type coils with air cores and a diameter of 1.8 m's out of PVC (Landler & Gollmann, 2011). The coils were placed parallel 0.9 m's apart from each other. Every coil had 21 turns of aluminum wire (section = 0.5 mm²) wrapped in plastic. We located the coils at the ground level and made a wooden structure of the base to avoid the use of metal. We measured the magnetic field strength with a 3-D flow valve (FLC3-70, Stefan Mayer Instruments) and we used a mechanical combustion generator (Dowel, 6.5 HP, Japan) as an energy source (220 V). The combustion generator on we placed the generator at 200 m distances to the sandbox to avoid additional stress to the individuals. The generator was kept on throughout the experiment. We employed a current limiter (VATRANS, Argentina) for generating a constant magnetic field in the coils (at 12 V, 50 A).

We carried out the tests in the circular arena and placed an adult amphibian in a release device in total darkness at the center of the sandbox. We constructed the release device using a black plastic container (0.2 × 0.2 m). We released the individuals after five minutes. The release device was removed from the arena using a pulley system, which allowed the operator to be far from the terrarium, not disturbing the experiment. As a directional response, we took the first contact of the individuals with the wall of the circular arena (Landler & Gollmann, 2011). Individuals, which could not reach the wall of the arena within five minutes, were excluded from the analysis.

We tested each individual twice, the first time under conditions of the Earth's Magnetic Field (EMF) and subsequently under the conditions of an added Induced Magnetic Field (IMF). To assess whether male *Rhinella arenarum* presents the "homing" behavior towards the reproductive site from which they were displaced, the displacement of each individual was first evaluated under the conditions of the Earth's magnetic field. Subsequently, to determine the ability of *R. arenarum* to perceive the Earth's magnetic field, the same individuals were evaluated under the conditions of an induced magnetic field (IMF). We followed the protocol described by Diego-Rasilla & Luengo (2002), where each test animal was used in the experiments consecutively. After testing each animal, we cleaned the sandbox with a paper towel and ethanol 50% to eliminate chemical substances produced by the tested individuals. To inverse, the horizontal component of the terrestrial magnetic field 180°, the Helmholtz type air-cored, coils were switched on. We returned the individuals to the ponds from which they were collected after the all experiment finished.

Experiments were carried out in as much darkness as possible (minimal generation of artificial light). Therefore, we obtained the records by a video camera with night vision (Sony Handycam, Japan), which was placed in the upper part of the circular arena (3.10 m height positions determined by the focus of the image) and which operated remotely from a monitor (Super Sonic Deluxe 5" Portable FC-9100 - USA) to reduce the stress of the animal to the maximum during the experiment.

We analyzed the data using standard circular statistics (Batschelet, 1981). We estimated mean vector length (r); mean direction (μ), circular variance, and concentration. We used Rayleigh's test to determine whether a distribution exhibited significant orientation. The V test was used to test closeness to expected orientation (i.e., home pond direction) (Batschelet, 1981; Mardia & Jupp, 1999). For the comparisons between the direction of toads between treatments, we used Watson-Williams F-Tests. All the tests had a confidence interval of 95%. The Wilcoxon

test was used to test differences in the time of response among treatments (Siegel, 1988). We performed all statistical analyses using the software Oriana version 4.02 of Kovach Computing Services and software Infostat (Di Rienzo et al., 2016).

RESULTS

The Earth's Magnetic Field in the study area at noon on September 23rd of 2012 had an average value of 0.21 ± 0.074 Gauss (range: $0.021 - 0.038$), and the Induced Magnetic Field at the center of the coils gave a maximal value of 0.45 ± 0.048 Gauss (range: 0.017 ± 0.098). From 44 studied anurans, only one individual, which did not move from the center of the arena, was excluded from the analyses. Under the influence of the Earth's Magnetic Field toads needed on average of 56.84 ± 35.56 s to touch the wall, and on average 48.49 ± 48.82 s, under an Induced Magnetic Field. No significant differences were found in the time of response among treatments (Wilcoxon test = -0.75 , $p = 0.45$, $n = 43$).

The direction of *R. arenarum* under the Earth's Magnetic Field was of $\mu = 305.41^\circ \pm 121.4^\circ$, of the vector mean (r) = 0.107 and of the concentration of data = 0.215 . The data exhibited a unimodal distribution (Test Rayleigh, $Z = 0.492$, $p = 0.61$). We not found significant differences in the orientation of *R. arenarum* respect of the spawning ponds (Test V (55.19°) = -0.036 , $p = 0.631$).

The direction of *R. arenarum* under an Induced magnetic field, but respect at North magnetic of the earth, was of $\mu = 224.58^\circ \pm 94.79^\circ$, of the vector mean (r) = 0.254 , and of the concentration of data = 0.526 . The data show a different tendency of the unimodal distribution, but not significant differences (Test Rayleigh, $Z = 2.784$, $p = 0.061$). The data show not significant differences respect of the spawning ponds (Test V (55.19°) = -0.25 , $p = 0.99$). Now, if we considered the North generated by the Induced Magnetic Field (at 180° to North Magnetic of the earth), the direction of *R. arenarum* was of $\mu = 44.58^\circ \pm 94.79^\circ$, of the vector mean (r) = 0.254 ,

and of the concentration of data = 0.526 . The data show a different orientation that deviates from a unimodal distribution, but it is statistically different respect of the spawning ponds (Test V (55.19°) = 0.25 , $p = 0.0092$).

The direction of the anurans aligned under the Earth's Magnetic Field and the spawning pond exhibited a unimodal distribution. In toads under the influence of the IMF showed a unimodal orientation that not coincided with the magnetic direction of the spawning ponds (Table 1). Concerning the magnitude of the mean vector and the concentration of the data, they were two times lower under EMF conditions than under IMF conditions (Table 1, Fig. 1) whereas the distribution patterns within the circular arena were significantly different among treatments ($F = 14.637$, $p = 0.0002$, $n = 43$ Watson-Williams F-Tests) and their orientation differed by 99° and 206° .

DISCUSSION

Our work studied the homing behavior of *Rhinella arenarum* towards the reproductive site and the influence of the magnetic field on their orientation to reorient their travel direction in response to an induced magnetic field were investigated. The results obtained demonstrate that, similar to other species of anurans, *R. arenarum* exhibits site fidelity (Roithmair, 1994; Neu et al., 2016; Nothacker et al., 2018). Ferguson et al. (1963) found that three species of anurans displayed homing behavior after returning to their site. Several studies show that male and female anurans can return to an initial capture site after experimental translocation (McVey et al., 1981; Crump 1986; Arcila-Pérez et al., 2020). The motivation for return might differ between sexes due to divergent selective pressures determining mating and reproductive success for males and females. The search motivation for territorial species' males is likely attributed to reclaiming a site more favorable for attracting mates than the release sites (Schlupp & Podloucky, 1994; Matthews, 2003).

Tabla 1. Estadística descriptiva para cada tratamiento; u (vector medio), r (longitud del vector medio), mediana, concentración, s (varianza circular), y s2 (desviación estándar circular). Test Rayleigh; $\alpha = 0.05$. V Test; CMT expectativa media 90° , CMI expectativa media 270° ; $\alpha = 0.05$. CMT: Campo magnético terrestre; CMI: Campo magnético inducido.

Table 1. Descriptive statistics for each treatment; u (mean vector), r (length of mean vector), median, concentration, s (circular variance), and s2 (circular standard deviation). Rayleigh Test; $\alpha = 0.05$. V Test; EMF expected mean 90° ; IMF expected mean 270° ; $\alpha = 0.05$. EMF: Earth magnetic field; IMF: Induced magnetic field.

Treatment	Descriptive statistics						Rayleigh Test		V Test		
	u	r	Concentration	Median	s	s2	Z	p	V	μ	p
EMF	3.573	0.107	0.216	332	0.893	121.07	1.012	0.363	0.007	0.089	0.465
IMF	220.408	0.265	0.556	232	0.735	93.36	6.183	0.002	0.172	2.279	0.011



In any case, males could return to the initial capture site due to the benefits associated with higher mating prospects. Perhaps a frog returns to its home spot because it is a good foraging site or perhaps a frog returns to its home spot because it is satisfactory, if not ideal, in terms of moisture, abundance of crevices in which to hide and sufficient overhanging vegetation to provide protection from the sun. Homing is clearly a strategy to minimize energy expenditure for locomotion, given the scarcity of suitable sites for reproduction, nutrition, and refuge.

Regarding the utilization of the amphibians' magnetic field for orienting to spawning sites, the results found in the experiments provided evidence that supports the idea that *R. arenarum* has many more orientation mechanisms than just the use of magnetic compass signals. However, a higher concentration in the orientation found in *R. arenarum* under the influence of the Induced Magnetic Field could demonstrate the importance of Earth's Magnetic Field, in the orientation in frogs (Fig. 1). Like

the results obtained by Landler & Gollmann (2011), Landler et al. (2016), Pail et al. (2020), in adult specimens of *Bufo bufo*, we obtained that adult specimens of *R. arenarum* tend to return to breeding pond after being moved from their site of origin. Our study reinforces the idea that the Earth's Magnetic Field could allow anuran specimens to reorientate their migration direction. The main difference between our work on *R. arenarum* and those carried out on *B. bufo* is that the collection of specimens in the former was carried out during migration, and in our study, they were collected in spawning ponds.

The ability of *R. arenarum* to orient itself to spawning ponds was consistent with results reported in several amphibian studies. Deutschlander et al. (1999) carried out the first experiences of the influence of the geomagnetic field on the orientation of *Notophthalmus viridescens* (Urodela) by studying the direction of these organisms by changing the magnetic field by a deviation of 90°. Diego-Rasilla et al. (2005) found in




Figura 1. Orientación de *R. arenarum* bajo el efecto del campo magnético terrestre (A) y el campo magnético inducido (B) relativo al norte geográfico. En cada diagrama circular, la dirección de las muestras, se agrupan en función del recorrido hasta el borde de la pista. Cada punto representa la orientación de un solo sapo. Las flechas representan los vectores medios de las distribuciones (el radio del círculo corresponde a un vector de longitud).

Figure 1. Orientation of *R. arenarum* under the effect of the Earth's Magnetic Field (A) and Induced Magnetic Field (B) relative to geographic north. In each circular diagram, the direction of the samples, are grouped concerning the course to the edge of the arena. Each dots represents the orientation of a single toad. The arrows represent the mean vectors of the distributions (radius of the circle corresponds to a vector length of 1).

Triturus alpestris that the magnetoreception mechanisms subject to celestial signals mainly explained the choice of direction toward the spawning pond. In another similar work carried out by Diego-Rasilla & Luengo (2007), they found that the *Lissotriton helveticus* was able to orient itself utilizing acoustic signals at short distances. However, at great distances from their spawning ponds, direction selection could only be done using the magnetic compass. Shakhparonov et al. (2017) studied adults of *Pelophylax ridibundus* and obtained similar results in comparison with the afore mentioned investigations. Diego-Rasilla et al. (2010) and Freake et al. (2002) proposed that the existing mechanisms of anurans and urodeles are traces of the evolution of sensory capacity in amphibians. Consequently, the sensitivity function of magnetoreception could have been present in a common ancestor of Lissamphibia, originating in the Lower Permian Period (Zhang & Wake, 2009). Despite all these studies on amphibian localization, it is not clear whether they can return home from unknown sites (Pail et al., 2020). It has been questioned whether true navigation represents a general ability of amphibians or if it could be a restricted phenomenon only present in a few species or even populations (Pasukonis et al., 2014; Sinch & Kirst, 2016).

The common toad, *R. arenarum* could orient itself to reproductive sites, being able to use the magnetic compass as a source of directional information. In our experiment, the *R. arenarum* specimens were shown to maintain the previous direction of migration in the “d” axis after being subjected to a field of inverse magnitude to the Earth’s Magnetic Field. However, its geographical location seems to depend more on the information on the map than on the integration of the route, to determine the initial location direction. The most likely source of information on the map is the geomagnetic field, which has been shown to play a critical role in the search process (Freake et al., 2002; 2006). However, various studies show the complexity of sensory integration in anurans. In distances less than 10 km it is very difficult to use a magnetic map (Komolkim et al., 2017). While olfactory signals could play a key role in orientation (Wallraf, 2004; Vignoli et al., 2012; Pail et al., 2020). Therefore, studies that complement the evaluation of various signals will serve to clarify the source of the orientation of anurans.

Despite finding a trend in our data and the lack of statistical support ($p = 0.063$), our results in this contribution provide the first direct in situ experimental support of the detection capacity of the terrestrial geomagnetic field in a species of toads in South America. Our research provides valuable insights into *R. arenarum* is homing behavior and their utilization of the magnetic field for orientation towards reproductive sites. The

species showcases remarkable site fidelity and demonstrates the intricate interplay of multiple orientation mechanisms, highlighting the complex nature of amphibian navigation. As in other amphibians, *R. arenarum* have a multisensory orientation and could use the terrestrial geomagnetic field as a navigation mechanism and location system to travel long distances to spawning ponds year after year. The orientation of amphibians is not a simple response to the environment but a complex process. We are confident that our work will enhance the understanding of amphibian orientation mechanisms. These findings contribute to a deeper understanding of anuran behavior and underscore the importance of Earth’s magnetic field in guiding their migrations and reproductive journeys. However, we believe that there is a need for a more comprehensive approach and a standardized methodology to identify the true mechanisms involved. Improved understanding will assist conservation efforts worldwide.

Acknowledgements.— We thank Heber Merenda, Yamil Rodriguez, Gonzalo Herrera, Daniela Moreno, Ariel Cataldo and Exequiel Gonzalez for their assistance in the field work. We especially thank to Barbara Vento, Jana Schön and Franco Lombino for the corrections in the translation and writing of the work in English. We thank the provincial fauna office of San Juan for permission SA y DS n° 1300-4736-2012 to conduct our research. This work was partially supported by The National Interuniversity Council of Argentina.

Ethical approval: All handling procedures were international guidelines and national and institutional regulations that apply to the care and use of animals.

CITED LITERATURE

- Arcila-Pérez, L.F., M.A. Atehortua-Vallejo & F. Vargas-Salinas. 2020. Homing in the rubí poison frog *Andinobates bombetes* (Dendrobatidae). *Copeia* 108:948-956.
- Batschelet, E. 1981. *Circular Statistics In Biology*. New York, NY. Academic Press Inc.
- Cabrera, A. 1971. Fitogeografía de la república argentina. *Boletín de la Sociedad Argentina de Botánica* 14:1-50.
- Camarillo-Naranjo, J.M., J.I. Álvarez-Francoso, N. Limones-Rodríguez, M.F. Pita-López & M. Aguilar-Alba. 2018. The global climate monitor system: from climate data-handling to knowledge dissemination. *International Journal of Digital Earth*. 8947:121.



- Crump, M.L. 1986. Homing and site fidelity in a neotropical frog, *Atelopus varius* (Bufonidae). *Copeia* 1986:438-444.
- Deutschlander, M.E., S.C. Borland & J.B. Phillips. 1999. Extraocular magnetic compass in newts. *Nature* 400:324-325.
- Di Rienzo, J., F. Casanoves, M. Balzarini, L. Gonzalez, M. Tablada & C. Robledo. 2016. InfoStat versión. Grupo InfoStat, FCA, Universidad Nacional de Córdoba, Argentina.
- Diego-Rasilla, F.J. & R.M. Luengo. 2007. Acoustic orientation in the palmate newt, *Lissotriton helveticus*. *Behavioral Ecology and Sociobiology* 61:1329-1335.
- Diego-Rasilla, F. J., R. M. Luengo & J. B. Phillips. 2008. Use of a magnetic compass for nocturnal homing orientation in the palmate newt, *Lissotriton helveticus*. *Ethology* 114:808-815.
- Diego-Rasilla, F.J. 2003. Homing ability and sensitivity to the geomagnetic field in the alpine newt, *Triturus alpestris*. *Ethology Ecology & Evolution* 15:251-259.
- Diego-Rasilla, F.J., R.M. Luengo & J.B. Phillips. 2005. Magnetic compass mediates nocturnal homing by the alpine newt, *Triturus alpestris*. *Behavioral Ecology and Sociobiology* 58:361-365.
- Diego-Rasilla, F.J., R.M. Luengo, J.B. Phillips. 2010. Light-dependent magnetic compass in Iberian green frog tadpoles. *Naturwissenschaften* 97:1077-1088.
- Diego-Rasilla, J. & R. Luengo. 2002. Celestial orientation in the marbled newt (*Triturus marmoratus*). *Journal of Ethology* 20:137-141.
- Diego-Rasilla, F. J., & Luengo, R. M. 2020. Magnetic compass orientation in common midwife toad tadpoles, *Alytes obstetricans* (Anura, Alytidae). *Journal of Ethology* 38:289-299.
- Endler, J. 1970. Kinesthetic orientation in the California newt (*Taricha torosa*). *Behaviour* 37:15-23.
- Ferguson, D. E. 1963. Orientation in three species of anuran amphibians. In *Orientierung der Tiere/Animal Orientation: Symposium in Garmisch-Partenkirchen* 17-21. 9. 1962. Pp. 128-134. Berlin, Heidelberg. Springer Berlin Heidelberg.
- Freake, M.J. & J.B. Phillips. 2005. Light-dependent shift in bullfrog tadpole magnetic compass orientation: evidence for a common magnetoreception mechanism in anuran and urodele amphibians. *Ethology* 111:241-254.
- Freake, M.J., C.S. Borland & J.B. Phillips. 2002. Use of a Magnetic Compass for Y-Axis Orientation in Larval Bullfrogs, *Rana catesbeiana*. *Copeia*. 2002:466-471.
- Freake, M.J., R. Muheim & J.B. Phillips. 2006. Magnetic maps in animals: a theory comes of age? *Quarterly Review of Biology* 81:327-347.
- Joly, P. & C. Miaud. 1993. How does a newt find its pond? The role of chemical cues in migrating newts (*Triturus alpestris*). *Ethology Ecology and Evolution* 5:447-455.
- Kramer, G. 1953. Wird die Sonnenhöhe bei der Heimfindeorientierung verwertet? *Journal für Ornithologie* 94:201-219.
- Landler L, D. Philippi, C. Nagl & G. Gollmann. 2016. Magnetic orientation in the common toad, *Bufo bufo*: a synopsis of arena experiments. *Mertensiella* 24:122-129.
- Landler, L. & G. Gollmann. 2011. Magnetic orientation of the Common Toad: establishing an arena approach for adult anurans. *Frontiers in Zoology* 8:1-6.
- Landler, L. 2022. Orientation and emigration of larval and juvenile amphibians: selected topics and hypotheses. *Amphibia-Reptilia* 43:1-11.
- Lohmann, K.J., C.M. Lohmann & N.F. Putman. 2007. Magnetic maps in animals: nature's GPS. *Journal of Experimental Biology* 210:3697-3705.
- Haddad, C.F. & C.P. Prado. 2005. Reproductive modes in frogs and their unexpected diversity in the Atlantic Forest of Brazil. *BioScience* 55:207-217.
- Rasilla, F J.D. & A.D. Luengo. 2022. A case of digital hindlimb malformations in "*Rana iberica*". *Boletín de la Asociación Herpetológica Española* 33:32-34.
- Mardia, K. & P. Jupp. 1999. *Directional Statistics*. Book Series. Wiley Series in Probability and Statistics. John Wiley & Sons, Inc.
- Matthews, K.R. 2003. Response of mountain yellow-legged frogs, *Rana muscosa*, to short distance translocation. *Journal of Herpetology* 3:621-626.



- McVey, M. E., R. G. Zahary, D. Perry & J. MacDougal. 1981. Territoriality and homing behavior in the poison dart frog (*Dendrobates pumilio*). *Copeia* 1981:1-8.
- Neu, C. P., S. S. Bisanz, J.A. Nothacker, M. Mayer & S. Lötters. 2016. Male and female home range behavior in the Neotropical poison frog *Ameerega trivittata* (Anura, Dendrobatidae) over two consecutive years. *South American Journal of Herpetology* 11:149-156.
- Nimpf, S. & D.A. Keays 2022. Myths in magnetosensation. *Iscience* 10:44-54.
- Nothacker, J. A., C.P. Neu, M. Mayer, N. Wagner & S. Lötters. 2018. Homing behavior in the Neotropical poison frog *Ameerega trivittata*. *Salamandra* 54:30-36.
- Pail, M., L. Landler & G. Gollmann. 2020. Orientation and navigation in *Bufo bufo*: a quest for repeatability of arena experiments. *Herpetozoa* 33:1-139.
- Pašukonis, A., M.C. Loretto, L. Landler, M. Ringler & W. Hödl. 2014. Homing trajectories and initial orientation in a Neotropical territorial frog, *Allobates femoralis* (Dendrobatidae). *Frontiers in Zoology* 11:1-9.
- Phillips, J.B. & S.C. Borland. 1992. Magnetic compassorientation is eliminated under near-infrared light in the eastern red-spotted newt *Notophthalmus viridescens*. *Animal Behaviour* 44:796-797.
- Phillips, J. B., K. Adler & S.C. Borland. 1995. True navigation by an amphibian. *Animal Behaviour* 50:855-858.
- Phillips, J.B. 1986. Magnetic compass orientation in the Eastern red-spotted newt (*Notophthalmus viridescens*). *Journal of Comparative Physiology A* 158:103-109.
- Phillips, J.B. 1996. Magnetic navigation. *Journal of Theoretical Biology* 180:309-319.
- Phillips, J.B. & F.J. Diego-Rasilla. 2022. The amphibian magnetic sense(s). *Journal of Comparative Physiology A* 208:723-742.
- Pittman, S.J. & C. A. McAlpine. 2003. Movements of marine fish and decapod crustaceans: process, theory and application. *Advances in Marine Biology* 44:205-294.
- Poblete, A. & M. Hryciw. 2017. Origen y variabilidad del recurso hídrico de los principales oasis de la Provincia de San Juan. *Boletín de Estudios Geográficos*. 107:9-20.
- Quinn, T. P. & A. H. Dittman 1990. Pacific salmon migrations and homing - mechanisms and adaptive significance. *Trends in Ecology & Evolution* 5:174-177.
- Quiroga L. & E. Sanabria 2012. Do reproductive parameters of *Rhinella arenarum* (Hensel, 1867) (Anura: Bufonidae) vary between the reproductive and postreproductive period? *Belgian Journal of Zoology* 142:68-73.
- Quiroga L.; E. Sanabria & J. Acosta 2004. Dimorfismo sexual en una población de *Bufo arenarum* (Anura: Bufonidae) en los Bañados de Zonda, San Juan, Argentina. *Boletín de la Sociedad Herpetológica Mexicana* 12:37-42.
- Quiroga L.; E. Sanabria & J.C. Acosta 2006. Size- and sex-dependent variation in diet of *Rhinella arenarum* (Anura: Bufonidae) in a wetland of San Juan, Argentina. *Journal of Herpetology* 43:311-317.
- Roithmair, M.E. 1994. Male territoriality and female mate selection in the dart-poison frog *Epipedobates trivittatus* (Dendrobatidae, Anura). *Copeia*, 1994:107-115.
- Sanabria E., L. Quiroga & A. Martino 2012. Variation in the thermoregulatory strategy of *Odontophrynus occidentalis* in the Monte desert, Argentina: response to the environmental constrain. *Journal of Experimental Zoology, part A. Ecological, Genetics and Physiology* 317:259-365.
- Sanabria E., L. Quiroga & J. Acosta. 2005. Patrones de actividad temporal estacional y uso de microhábitat de una población de adultos de *Bufo arenarum*, en los humedales de Zonda, San Juan, Argentina. *Boletín de la Sociedad Herpetológica Mexicana* 13:61-65.
- Sanchez, L. C., R.C. Lajmanovich, P. Peltzer, A.S. Manzano, C.M. Junges, & A.M. Attademo. 2014. First evidence of the effects of agricultural activities on gonadal form and function in *Rhinella fernandezae* and *Dendropsophus sanborni* (Amphibia: Anura) from Entre Ríos Province, Argentina. *Acta Herpetologica* 9:75-88.
- Schlupp I. & R. Podlucky. 1994. Changes in breeding site fidelity: a combined study of conservation and behaviour in the common toad *Bufo bufo*. *Biological Conservation* 69:285-291.



- Shakhparonov, V.V., A.P. Golovlev, E.E. Grytshyshina & A.A. Bolshakova. 2022. Orientation in the European common frog *Rana temporaria* during the first wintering migration. *Journal Experimental Biology* 225:jeb243761.
- Shakhparonov, V.V. & S.V. Ogurtsov. 2017. Marsh frogs, *Pelophylax ridibundus*, determine migratory direction by magnetic field. *Journal of Comparative Physiology A* 203:35-43.
- Siegel, S. & N.J. Castellan. 1988. *Nonparametric Statistics for the Behavioural Sciences*. McGraw-Hill, New York, USA.
- Sinsch, U. 2014. Movement ecology of amphibians: from individual migratory behaviour to spatially structured populations in heterogeneous landscapes. *Canadian Journal of Zoology* 92:491-502.
- Sinsch, U. 1990. The orientation behaviour of three toad species (genus *Bufo*) displaced from the breeding site. in *Biology and Physiology of Amphibians*. 38:75-83. *Fortschritte der Zoologie*. New York.
- Sinsch, U. & C. Kirst. 2016. Homeward orientation of displaced newts (*Triturus cristatus*, *Lissotriton vulgaris*) is restricted to the range of routine movements. *Ethology Ecology & Evolution* 28:312-328.
- Skiles, D.D. 1985. The geomagnetic field its nature, history, and biological relevance. In *Magnetite Biomineralization and Magnetoreception in Organisms*. In Kirschvink, J.L. (Ed.). Springer, Boston, MA.
- Sotomayor, V., C. Lascano, A.M.P. de D'Angelo & A. Venturino. 2012. Developmental and polyamine metabolism alterations in *Rhinella arenarum* embryos exposed to the organophosphate chlorpyrifos. *Environmental Toxicology and Chemistry* 31:2052-2058.
- Victoria, J.A. 1999. Simulación matemática del sistema embalse de Ullum-Cuenca de agua subterránea. Instituto Nacional de Agua y Ambiente, San Juan, Argentina.
- Vignoli, L., R. Silici, A.M. Bissattini & M.A. Bologna. 2012. Aspects of olfactory mediated orientation and communication in *Salamandrina perspicillata* (Amphibia Caudata): an experimental approach. *Ethology Ecology & Evolution* 24:165-173.
- Walcott, C. 1996. Pigeon homing: observations, experiments and confusions. *Journal of Experimental Biology* 199:21-27.
- Wallraff, H. G. 2004. Avian olfactory navigation: its empirical foundation and conceptual state. *Animal Behaviour* 67:189-204.
- Wells, K. D. 2007. *The Ecology and Behavior of Amphibians*. University of Chicago Press, Chicago, USA.
- Wiltschko, R. & W. Wiltschko. 2012. Magnetic Orientation in Animals. In *Magnetic Orientation in Animals*, Vol. 33. Springer Science & Business Media, Heidelberg, Germany.
- Young, B.E., S.N. Stuart, J.S. Chanson, N.A. Cox & M.T. Boucher. 2004. *Joyas que Están Desapareciendo: El Estado de los Anfibios en el Nuevo*. NatureServe, Arlington, Virginia.
- Zhang, P. & D.B. Wake. 2009. Higher-level salamander relationships and divergence dates inferred from complete mitochondrial genomes. *Molecular Phylogenetics and Evolution* 53:492-508.

